

p -adic CCR: Galois group representations, cyclic dynamics, and zeta-functions

Andreas Martin Lisewski*
Belgradstrasse 19, 80796 Munich, Germany

We consider a model of cyclic time evolution for Kochubei's p -adic realization of the canonical commutation relations (CCR). Connections to Kubota-Leopoldt p -adic zeta-functions and to arithmetic quantum theories such as the Bost-Connes model are examined.

I. INTRODUCTION

The physical interpretation of arithmetic quantum theories has gained attention in recent years. Briefly, these quantum theories are those which exhibit Dirichlet L -series as thermal partition functions in their statistical formulation. Grounded on mathematical work of Julia [1] and Bost-Connes [2], it was realized that there exist similarities to certain aspects of string theory at finite temperature, or to other statistical systems such as the two-dimensional Ising model. For example, the pole of the Riemann ζ -function $\zeta(\beta)$ at $\beta = 1$ may be understood as an indicator of a phase transition at Hagedorn temperature $\beta^{-1} = 1$ [1, 3], and the functional equation for the ζ -function may be conceived as a duality relation in analogy to the Kramers-Wannier relation for the two-dimensional Ising model [1]. However, arithmetic quantum theories have not been treated as truly physical theories so far, which in other words means that until now none of them has been identified as a realistic model of a physical system.

The motivation of this note is the question whether there are p -adic quantum systems that share certain aspects of arithmetic quantum theories. In contrast to conventional quantum theory, p -adic quantum systems such as the one investigated here are realized on spaces over the field $\mathbb{Q}_p$ of p -adic numbers (p -adic fields, which in addition are algebraically closed are not regarded here.). Our main result is that the probably most elementary p -adic quantum system, namely the quantum harmonic oscillator whose generators satisfy the canonical commutation relations (CCR), exhibits a structural similarity to arithmetic quantum theories. Based on Kochubei's p -adic construction of the CCR [4], we show that the Galois group $\text{Gal}(\mathbb{Q}_p/\mathbb{Q})$ of the abelian extension $\mathbb{Q}_p$ of the rational numbers $\mathbb{Q}$ naturally leads to a representation ρ on a one-dimensional vector space containing the oscillator's ground state. This leads us to a proposition stating that there exists a group homomorphism from the Galois group $\text{Gal}(\mathbb{Q}_p/\mathbb{Q})$ to the group of automorphisms acting on the algebra of bounded linear operators over the Banach space of continuous functions on $\mathbb{Z}_p$. From the perspective of C^* -dynamical systems our representation motivates us to regard $\text{Gal}(\mathbb{Q}_p/\mathbb{Q})$ as the one-parameter group of cyclic time evolution for the p -adic quantum sys-

tem. Further, the one-dimensional representation itself turns out to be directly related to an Artin p -adic L -function associated to ρ . In our case the latter is equal to the Kubota-Leopoldt p -adic ζ -function. We close with comments on the physical implications of our results and indicate obvious connections to the Bost-Connes arithmetic quantum system.

II. p -ADIC CCR

We follow the work of Kochubei [4]. Let p be a prime number, $\mathbb{Q}_p$ the field of p -adic numbers, $\mathbb{Z}_p$ the ring of p -adic integers. Denote $C(\mathbb{Z}_p, \mathbb{Q}_p)$ the Banach space of $\mathbb{Q}_p$ -valued continuous functions on $\mathbb{Z}_p$ equipped with sup-norm. For $n \geq 1$ the functions

$$P_n(x) = \frac{x(x-1) \cdots (x-n+1)}{n!} \quad (1)$$

with $P_0(x) := 1$ form complete orthonormal system, i.e. the Mahler basis, of $C(\mathbb{Z}_p, \mathbb{Q}_p)$. Thus every function $f \in C(\mathbb{Z}_p, \mathbb{Q}_p)$ admits a unique expansion

$$f(x) = \sum_{n=0}^{\infty} c_n P_n(x), \quad c_n \in \mathbb{Q}_p \quad (2)$$

with $|c_n|_p \rightarrow 0$, and $\|f\|_p = \sup_n |c_n|_p$, and with $|\cdot|_p$ being the absolute value of a p -adic number. Consider now the following operators on $C(\mathbb{Z}_p, \mathbb{Q}_p)$:

$$\begin{aligned} (a^+ f)(x) &= x f(x-1) \\ (a^- f)(x) &= f(x+1) - f(x). \end{aligned} \quad (3)$$

This construction is well defined as it directly follows from the definition of the $\|\cdot\|_p$ norm that $a^{\pm}$ are bounded operators. A direct calculation of the commutator gives the CCR

$$[a^-, a^+] = 1. \quad (4)$$

Moreover, we obtain

$$\begin{aligned} a^- P_n &= P_{n-1}, \quad n \geq 1, \quad a^- P_0 = 0 \\ a^+ P_n &= (n+1) P_{n+1}, \quad n \geq 0 \end{aligned}$$

thus a^+, a^- are clearly analogues of the creation and annihilation operators. Setting $H = a^+ a^-$ we obtain $HP_n = nP_n$, and $[H, a^{\pm}] = \pm a^{\pm}$. The operator H has a complete system of eigenvectors and its discrete spectrum coincides with the set of non-negative integers

*Electronic address: martin.lisewski@soundate.com

Z_+ . The whole spectrum of H , however, equals Z_p , that is the $|\cdot|_p$ -closure of Z_+ in Q_p . The density of Z_+ in Z_p implies that the kernel of a^- consists of constant functions on Z_p . This fact may be used to show that a^-, a^+ form an irreducible representation of the CCR. A remarkable property is that every Banach space over Q_p having an infinite countable orthonormal basis is isomorphic to $C(Z_p, Q_p)$.

III. GROUND STATE STRUCTURE AND GALOIS GROUP REPRESENTATIONS

As ground state of our p -adic quantum mechanical system we denote any continuous function $\Omega \in C(Z_p, Q_p)$ with $a^-\Omega = 0$ and $\Omega|_{Z_p} = 1$. If we enlarge the domain of ground states and consider continuous functions $\Omega : X \rightarrow Q_p \in C(X, Q_p)$, $Z_p \subset X$, then we may expect more than just the trivial case, i.e. a globally constant mapping, because now any locally constant function on Z_p with $\Omega|_{Z_p} = 1$ may be regarded as a ground state, too. It is then convenient to look at the equivalence class Ω of such ground states defined upon the relation

$$\Omega_1 \sim \Omega_2 \Leftrightarrow \Omega_1, \Omega_2 \in C(X, Q_p) : \Omega_1|_{Z_p} = \Omega_2|_{Z_p} = 1.$$

As Q_p is a field there are $p-1$ primitive roots of unity $\mu^{p-1} = \{\zeta_1, \dots, \zeta_{p-1}\}$, $\zeta_j \in Z_p$ for all $j \in [1, p-1]$, together being the only roots of $x^{p-1} - 1$ in Q_p . These roots are the image of the Teichmüller character ω , $\omega : Z_p^* \rightarrow \mu^{p-1}$, with the representation property $\omega(\alpha)\omega(\beta) = \omega(\alpha\beta)$, $\alpha, \beta \in Z_p^*$. Since Q_p is an extension field of the field of the rational numbers Q , one has the corresponding Galois group $\text{Gal}(Q_p/Q) \simeq Z_p^*$. Consider the one-dimensional Q_p -vector space $V_0 = \{c\Omega : c \in Q_p\}$. Then, for every fixed $\kappa_0 \in \{0, \dots, p-2\}$ the Teichmüller character ω admits a faithful one-dimensional representation $\rho_{\kappa_0} : \text{Gal}(Q_p/Q) \rightarrow \text{Aut}(V_0)$ of the Galois group, viz.

$$\rho_{\kappa_0}(\alpha)\Omega = \omega(\alpha)^{\kappa_0}\Omega, \quad \forall \alpha \in Z_p^*. \quad (5)$$

We stress that ρ_{κ_0} is a continuous representation, since ω extends to a (uniformly) continuous function $\omega \in C(Z_p, Q_p)$. Since $Z_p^* \simeq Z_{p-1}$, equation (5) may also be written as

$$\rho_{\kappa_0}(\alpha)\Omega = \zeta_{p-1}^{\kappa_0 t(\alpha)}\Omega, \quad (6)$$

for a suitably chosen $t(\alpha) \in \{0, \dots, p-2\}$ and ζ_{p-1} is the primitive root of unity of degree $p-1$.

Let $B(C(Z_p, Q_p)) \equiv B$ be the algebra over Q_p of bounded linear operators on $C(Z_p, Q_p)$, i.e. $B = \{A \in B : \|Af\| < \infty, \forall f \in C(Z_p, Q_p)\}$. Our next task is to show that the above representation ρ naturally induces a representation $\rho' : \text{Gal}(Q_p/Q) \rightarrow \text{Aut}(B)$. Every $A \in B$ may uniquely be represented as a set A' of ordered pairs: $A' = \{(f, Af) : f \in C(Z_p, Q_p)\}$. Obviously, the collection of all such sets forms an algebra $B' \simeq B$. We introduce the *van der Put basis* $\{e_n : n \in N_0\}$ of $C(Z_p, Q_p)$ as follows:

$e_0 = 1$ and for $n > 0$, e_n is the characteristic function of the disc $D_n = \{x \in Z_p : |x - n|_p < 1/n\}$. Then for every $f \in C(Z_p, Q_p)$ one obtains the uniformly convergent series:

$$f(x) = \sum_{n=0}^{\infty} v_n e_n(x)$$

with $v_0 = f(0)$ and $v_n = f(n) - f(n_-)$; n_- is defined through the Hensel expansion given for any $n \in N$: $n = n_0 + n_1p + \dots + n_sp^s$ with $n_s \neq 0$. Then $n_- = n_0 + n_1p + \dots + n_{s-1}p^{s-1}$. Thus for every $A' \in B'$ and every $f \in C(Z_p, Q_p)$ we have

$$(f, Af) = \left(\sum_{n=0}^{\infty} v_n(f) e_n(x), \sum_{n=0}^{\infty} v_n(Af) e_n(x) \right). \quad (7)$$

But every e_n is locally constant on Z_p ; this is because N is dense in Z_p and each disc D_n exhausts all the elements of Z_p . Hence, we may write

$$(f, Af) = \left(\sum_{n=0}^{\infty} v_n(f) \Omega, \sum_{n=0}^{\infty} v_n(Af) \Omega \right). \quad (8)$$

As the expansion in the van der Put basis is uniformly convergent, we have the following mapping due to equation (6)

$$\rho'_{\kappa_0} : t \mapsto \left(\sum_{n=0}^{\infty} v_n(f) \Omega, \sum_{n=0}^{\infty} v_n(Af) \zeta_{p-1}^{\kappa_0 t} \Omega \right) \quad (9)$$

with $t \in Z_{p-1}$. Since ρ_{κ_0} is faithful this mapping determines an automorphism on B' , and so it does on B . Thus in summary, we have the proposition

Proposition III.1. *Let the tuple $(C(Z_p, Q_p), a^{\pm}, B)$ be the p -adic quantum mechanical system as previously defined. Let further $\Omega \in V_0$ denote the ground state of this system. Then for every $\kappa_0 \in Z_{p-1}$ there is a faithful representation $\rho'_{\kappa_0} : \text{Gal}(Q_p/Q) \rightarrow \text{Aut}(B)$ uniquely determined by the one-dimensional continuous representation $\rho_{\kappa_0} : \text{Gal}(Q_p/Q) \rightarrow \text{Aut}(V_0)$.*

IV. p -ADIC ζ -FUNCTIONS

Finite dimensional representations of Galois groups associated to field extensions exhibit a remarkable relation to certain L -functions. In particular, due to the pioneering work of Deligne and Ribet [5] one can naturally define a p -adic L -function $L_p(s, \rho)$ to any even representation ρ on a vector space over a p -adic field such as is Q_p . This function is referred to as p -adic Artin L -function associated to ρ . Let C_p be the smallest extension field of Q_p that is algebraically closed and complete with respect to $|\cdot|_p$. Then a one-dimensional representation of $\text{Gal}(C_p/Q)$ generally reads as

$$\rho(\sigma_{\alpha})e = \chi(\alpha)e$$

where e is the basis of vector space $V = C_p e$. Here, ρ is even when $\chi(-1) = 1$. For even representations of this kind the p -adic Artin L -function associated to ρ becomes the p -adic Dirichlet L -function originally introduced by Kubota and Leopoldt [6]. If we now restrict our view to extension field Q_p then the one-dimensional representations of $\text{Gal}(Q_p/Q)$ are those given by equation (5) and - equivalently - by equation (6) with even characters $\chi = \omega^{\kappa_0}$. In this case the Kubota-Leopoldt p -adic Dirichlet L -function is defined through a Q_p -valued measure $\mu(x)$ and it reads (For an introduction to p -adic integration and p -adic L -functions, see [7].)

$$\zeta_{p,\kappa_0}(s) = \frac{1}{\langle r \rangle^{1-s} \omega(r)^{\kappa_0} - 1} \int_{Z_p^*} \langle x \rangle^{-s} \omega(x)^{\kappa_0-1} d\mu(x) \quad (10)$$

where r is any integer prime to p , $\langle r \rangle = r/\omega$, $\langle x \rangle = x/\omega$, and the parameter $\kappa_0 \in \{0, \dots, p-2\}$ depicts the *branches* of the p -adic ζ -function. p -adic Dirichlet L -functions have the remarkable property that they interpolate complex Dirichlet L -functions at algebraic values, i.e. values for $1-k$ with $k \in N_0$. So, for example, if we choose $p=2$ then we have

$$\zeta_{2,0}(1-k) = (1-2^{k-1}) \zeta(1-k) = (2^{k-1}-1) \frac{B_k}{k},$$

where ζ is the Riemann ζ -function and B_k is the k th Bernoulli number. Despite this close analogy to complex Dirichlet L -functions, p -adic Dirichlet L -functions are less understood than their complex counterparts. For instance, a general functional equation for p -adic Dirichlet L -functions is unknown, and their values for natural numbers larger than one are not known either.

V. DISCUSSION

Proposition III.1 plays a similar role for our p -adic quantum mechanical system as does a proper group homeomorphism $\alpha : R \rightarrow \text{Aut}(\mathcal{A})$ play for some C^* -dynamical system within the framework of conventional quantum mechanics, i.e. the role of time evolution within the algebra of observables. In the latter, $\mathcal{A}$ usually is the C^* -algebra of bounded linear operators on some complex Hilbert space.

Despite this obvious similarity there are significant differences. Recall that Q_p is not an algebraically closed field; for example, the equation $x^2 + 1 = 0$ does not have a solution in Q_p . Therefore, our construction shows that with p -adic analysis over Q_p it is possible to represent a non-trivial quantum mechanical system and its time evolution on a real (in the field-theoretic sense) one-dimensional vector space- something that is totally missing for one-dimensional (Hilbert) spaces over R . In fact, if we try to apply our results to the case of real numbers and consequently identify $Q_\infty = R$ then the corresponding Galois group becomes $\text{Gal}(R/Q) \simeq \{\text{id}\}$, i.e. the trivial group. Now the only roots of unity in R are $\{\pm 1\}$. Since there is no continuous homeomorphism $\rho : \{\text{id}\} \rightarrow \{\pm 1\}$, there is no non-trivial time evolution either. For the p -adic case, on the other hand, we have a cyclic time structure defined for each branch that is depicted by a value of κ_0 . Surely, the physical meaning of a p -adic time parameter $t \in Z_{p-1}$ is far from being clear, because the usual complete ordering of real numbers is lost here. The latter property, however, has always been considered as being essential for a physical time parameter. So, p -adic quantum systems such as the one investigated here may indicate that R does not give the only possible parametrization of time.

Finally, we want to briefly mention an obvious relation between the p -adic quantum system as introduced in this note and the arithmetic quantum system of Bost and Connes. In both systems the Galois group of an abelian extension of Q has been represented on the corresponding algebra of observables. And in both systems a Dirichlet L -function naturally occurs as a consequence. In the Bost-Connes system this L -function is the Riemann ζ -function $\zeta(\beta)$ which at the same time turns out to be the partition function of the quantum statistical system at temperature β^{-1} . Therefore, one may wonder whether it is admissible to regard $\zeta_{p,\kappa_0}(s)$ as a p -adic partition function, and whether s may become the inverse p -adic temperature in any reasonable sense. Further, it is interesting to observe that our p -adic quantum system possesses an extra symmetry not present in the Bost-Connes system: the one-parameter group representation of $\text{Gal}(Q_p/Q)$ appears twice as realized through the parameters κ_0 and t . Thus $\text{Gal}(Q_p/Q)$ plays a double-role here in the sense that it is a symmetry group of the quantum system as well as it is the system's dynamical group representing time.

-
- [1] B. Julia, J. Phys. France **50**, 1371 (1989).
 - [2] J.-B. Bost and A. Connes, Selecta Mathematica **1**, 411 (1995).
 - [3] D. Spector, J. Math. Phys. **39**, 1919 (1998).
 - [4] A. N. Kochubei, J. Phys. A **29**, 6375 (1996).
 - [5] P. Deligne and K. Ribet (1979), IHES preprint/October

- 1979.
- [6] T. Kubota and H. W. Leopoldt, J. Reine und Angew. Math. **214/215**, 328 (1964).
- [7] N. Koblitz, *p-adic Analysis: a Short Course on Recent Work* (Cambridge University Press, 1980).